

On certain $\mathcal{C}$ -ultrahomogeneous graphs obtained from cubic distance transitive graphs

Italo J. Dejter
University of Puerto Rico
Rio Piedras, PR 00931-3355
ijdejter@uprrp.edu

Abstract

The notion of a $\mathcal{C}$ -ultrahomogeneous (or $\mathcal{C}$ -UH) graph due to D. Isaksen et al. is adapted for digraphs and applied to the cubic distance-transitive graphs, considered both as graphs and digraphs, when $\mathcal{C}$ is formed by shortest cycles and $(k-1)$ -paths, with $k = \text{arc-transitivity}$. Moreover, $(k-1)$ -powers of shortest cycles taken with orientation assignments that make these graphs become $\mathcal{C}$ -UH digraphs are ‘zipped’ into novel $\mathcal{C}$ -UH graphs, with $\mathcal{C}$ formed by copies of K_3 , K_4 , C_7 and $L(Q_3)$.

Keywords: ultrahomogeneous graph; digraph; shortest cycle; arc-transitivity

1 Preliminaries

The study of ultrahomogeneous graphs (resp. digraphs) can be traced back to [16], [10], [15], [3] and [12], (resp. [9], [14] and [4]). Following a line of research initiated by [13], given a collection $\mathcal{C}$ of (di)graphs closed under isomorphisms, a (di)graph G is said to be $\mathcal{C}$ -ultrahomogeneous (or $\mathcal{C}$ -UH) if every isomorphism between two induced members of $\mathcal{C}$ in G extends to an automorphism of G . If $\mathcal{C} = \{H\}$ is the isomorphism class of a graph H , we say that such a G is $\{H\}$ -UH or H -UH. In [13], $\mathcal{C}$ -UH graphs are defined and studied when $\mathcal{C}$ is the collection of either **(a)** the complete graphs, or **(b)** the disjoint unions of complete graphs, or **(c)** the complements of those unions. In [6], a $\{K_4, K_{2,2,2}\}$ -UH graph X that fastens objects of (a) and (c), namely K_4 and $K_{2,2,2}$, is presented.

We may consider a graph G as a digraph by considering each edge e of G as a pair of oppositely oriented (or O-O) arcs $\vec{e}$ and $(\vec{e})^{-1}$. Then, ‘zipping’ $\vec{e}$ and $(\vec{e})^{-1}$ allows to recover e , a technique to be used below. In [7], however, a strongly connected C_4 -UH oriented graph without O-O arcs was presented.

Let M be a sub(di)graph of a (di)graph H and let G be both an M -UH and an H -UH (di)graph. We say that G is a (fastened) $(H; M)$ -UH (di)graph if, given a copy H_0 of H in G containing a copy M_0 of M , then there exists exactly one copy $H_1 \neq H_0$ of H in G with $V(H_0) \cap V(H_1) = V(M_0)$ and $A(H_0) \cap \bar{A}(H_1) = A(M_0)$, where $\bar{A}(H_1)$ is formed by those arcs $(\vec{e})^{-1}$ whose orientations are reversed with

respect to those of the arcs $\vec{e}$ of $A(H_1)$, and moreover: no more vertices or arcs than those in M_0 are shared by H_0 and H_1 . The directed case here is used in the constructions of Section 3. In the undirected case, the vertex and arc conditions above can be condensed as $H_0 \cap H_1 = M_0$; this is generalized by saying that an $(H; M)$ -UH graph G is an ℓ -fastened $(H; M)$ -UH graph if given a copy H_0 of H in G containing a copy M_0 of M , then there exist exactly ℓ copies $H_i \neq H_0$ of H in G such that $H_i \cap H_0 = M_0$, for each one of $i = 1, 2, \dots, \ell$, and such that no more vertices or edges than those in M_0 are shared by each two of $H_0, H_1, \dots, H_\ell$.

To better state the cited result from [6], let H_i be a connected graph, for $i = 1, 2$, with $H_i \not\subset H_j$, ($i \neq j \in \{1, 2\}$). A graph G is a $(K_2$ -fastened) $\{H_2, H_1\}_{K_2}$ -UH graph if for each $i \in \{1, 2\}$: **(a)** G is an H_i -UH graph; **(b)** G is representable as an edge-disjoint union of a number n_i of induced copies of H_i ; **(c)** G has a constant number m_i of copies of H_i incident at each vertex, with no two such copies sharing more than one vertex; **(d)** G has exactly n_i copies of H_i as induced subgraphs; **(e)** G has each edge in exactly one copy of H_i .

In [6], it is shown that X is a $\{K_4, K_{2,2,2}\}_{K_2}$ -UH graph. This is the first known such $\mathcal{C}$ -UH graph that is not a line graph, a fact relevant in view, for example, of the line graphs of the d -cubes, $3 \leq d \in \mathbf{Z}$, (say the cuboctahedron $L(Q_3)$), which are $\{K_d, K_{2,2}\}_{K_2}$ -UH. In addition, X has exactly 42 copies of K_4 and 21 copies of $K_{2,2,2}$ as subgraphs. X also has four copies of K_4 and three copies of $K_{2,2,2}$ incident at each one of its 42 vertices. Considering self-dual configurations and their Levi and Menger graphs as in [5], X is seen to be the Menger graph of a self-dual (42_4) -configuration.

We will work with the cubic distance-transitive (or CDT) graphs G , [2]:

CDT graph G	n	d	g	k	η	a	b	h	κ
Tetrahedral graph K_4	4	1	3	2	4	24	0	1	1
Thomsen graph $K_{3,3}$	6	2	4	3	9	72	1	1	2
3-cube graph Q_3	8	3	4	2	6	48	1	1	1
Petersen graph	10	2	5	3	12	120	0	0	0
Heawood graph	14	3	6	4	28	336	1	1	0
Pappus graph	18	4	6	3	18	216	1	1	3
Dodecahedral graph	20	5	5	2	12	120	0	1	1
Desargues graph	20	5	6	3	20	240	1	1	3
Coxeter graph	28	4	7	3	24	336	0	0	3
Tutte 8-cage	30	4	8	5	90	1440	1	1	2
Foster graph	90	8	10	5	216	4320	1	1	0
Biggs-Smith graph	102	7	9	4	136	2448	0	1	3

where n, d, g, k, η and a are order, diameter, girth, AT or arc-transitivity, number of g -cycles and number of automorphisms, respectively, with b (resp. h) = 1 if G is bipartite (resp. hamiltonian) and = 0 otherwise, and κ defined as follows: let P_k and $\vec{P}_k$ be respectively a $(k-1)$ -path and a directed $(k-1)$ -path (of length $k-1$); let C_g and $\vec{C}_g$ be respectively a cycle and a directed cycle of length g ; then (see Theorem 2 below): $\kappa = 0$, if G is not $(\vec{C}_g; \vec{P}_k)$ -UH; $\kappa = 1$, if G is planar; $\kappa = 2$, if G is $(\vec{C}_g; \vec{P}_k)$ -UH with $g = 2(k-1)$; $\kappa = 3$, if G is $(\vec{C}_g; \vec{P}_k)$ -UH with $g > 2(k-1)$.

Given a finite graph H and a subgraph M of H with $|V(H)| > 3$, we say that a graph G is (strongly fastened) SF $(H; M)$ -UH if there is a descending sequence of connected subgraphs $M = M_1, \dots, M_{|V(H)|-2} \equiv K_2$ such that: **(a)**

M_{i+1} is obtained from M_i by the deletion of a vertex, for $i = 1, \dots, |V(H)| - 3$ and **(b)** G is a $(2^i - 1)$ -fastened $(H; M_i)$ -UH graph, for $i = 1, \dots, |V(H)| - 2$. Theorem 1 below asserts that every CDT graph is an SF $(C_g; P_k)$ -UH graph.

Given a graph C and $0 < k \in \mathbf{Z}$ such that k is at most the diameter of C , recall that the k -power graph C^k of C has $V(C^k) = V(C)$ and that two vertices are adjacent in C^k if and only if they are at distance k in C . Theorem 2 establishes which CDT graphs are $(\vec{C}_g; \vec{P}_k)$ -UH digraphs. Elevating the resulting oriented cycles to the $(k-1)$ -power enables the construction, in Section 3 and in [8], of fastened $\mathcal{C}$ -UH graphs, with $\mathcal{C}$ formed by copies of K_3 , K_4 , C_7 and $L(Q_3)$, when $\kappa = 3$, via ‘zipping’ of the O-O induced $(k-1)$ -arcs shared (as $(k-1)$ -paths) by pairs of O-O g -cycles. In particular: **(a)** the Pappus (resp. Desargues) graph yields the disjoint union of two copies of the Menger graph of the self-dual (9_3) - (resp. (10_3) -) configuration, [5]; **(b)** the Coxeter graph yields the Klein graph on 56 vertices, [1, 17]; **(c)** the Biggs-Smith graph yields the Menger graph of a self-dual (102_4) -configuration, a non-line-graphical $\{K_4, L(Q_3)\}_{K_3}$ -UH graph, (in contrast with the self-dual (42_4) -configuration of [6], whose Menger graph X is $\{K_4, K_{2,2,2}\}_{K_2}$ -UH), treated separately in [8].

2 (C_g, P_k) -UH properties of CDT graphs

Theorem 1 *Let G be a CDT graph of girth $= g$ and $AT = k$. Then G is an SF $(C_g; P_{i+2})$ -UH graph, for $i = 0, 1, \dots, k - 2$. In particular, G is a $(C_g; P_k)$ -UH graph and has exactly $2^{k-2}3ng^{-1}$ g -cycles.*

Proof. We have to see that each CDT graph G with girth $= g$ and $AT = k$ is a $(2^i - 1)$ -fastened $(C_g; P_{k-i})$ -UH graph, for $i = 0, 1, \dots, k - 2$. In fact, each $(k - i - 1)$ -path $P = P_{k-i}$ of any such G is shared exactly by 2^i g -cycles of G , for $i = 0, 1, \dots, k - 2$. Moreover, each two of these 2^i g -cycles have just P in common. This and a simple counting argument for the number of g -cycles, as cited in the table above, yield the assertions in the statement. $\square$

Theorem 2 *The CDT graphs G of girth $= g$ and $AT = k$ that are not $(\vec{C}_g; \vec{P}_k)$ -UH digraphs are the Petersen graph, the Heawood graph and the Foster graph. The remaining nine CDT graphs are $(\vec{C}_g; \vec{P}_k)$ -UH.*

Proof. Given a $(\vec{C}_g; \vec{P}_k)$ -UH graph G , an assignment of an orientation to each g -cycle of G such that the two g -cycles shared by each $(k - 1)$ -path receive opposite orientations yields a $(\vec{C}_g; \vec{P}_k)$ -orientation assignment (or $(\vec{C}_g; \vec{P}_k)$ -OA). The collection of η oriented g -cycles corresponding to the η g -cycles of G , for a particular $(\vec{C}_g; \vec{P}_k)$ -OA will be called an $(\eta\vec{C}_g; \vec{P}_k)$ -OAC.

The graph $G = K_4$ on vertex set $\{1, 2, 3, 0\}$ admits the $(4\vec{C}_3; \vec{P}_2)$ -OAC $\{(123), (210), (301), (032)\}$. The graph $G = K_{3,3}$ obtained from K_6 (with vertex set $\{1, 2, 3, 4, 5, 0\}$) by deleting the edges of the triangles $(1, 3, 5)$ and $(2, 4, 0)$ admits the $(9\vec{C}_4; \vec{P}_3)$ -OAC $\{(1234), (3210), (4325), (1430), (2145), (0125), (5230),$

(0345), (5410)). The graph $G = Q_3$ with vertex set $\{0, \dots, 7\}$ and edge set $\{01, 23, 45, 67, 02, 13, 46, 57, 04, 15, 26, 37\}$ admits the $(6\vec{C}_4; \vec{P}_2)$ -OAC $\{(0132), (1045), (3157), (2376), (0264), (4675)\}$.

If $G = Pet$ is the Petersen graph, then G can be obtained from the disjoint union of the 5-cycles $\mu_\infty = (u_0u_1u_2u_3u_4)$ and $\nu_\infty = (v_0v_2v_4v_1v_3)$ by the addition of the edges (u_x, v_x) , for $x \in \mathbf{Z}_5$. Apart from the two 5-cycles given above, the other ten 5-cycles of G can be denoted by $\mu_x = (u_{x-1}u_xu_{x+1}v_{x+1}v_{x-1})$ and $\nu_x = (v_{x-2}v_xv_{x+2}u_{x+2}u_{x-2})$, for each $x \in \mathbf{Z}_5$. Then the following sequence of alternating 6-cycles and 2-arcs starts and ends up with opposite orientations: $\mu_2^-(u_3u_2u_1)\mu_\infty^+(u_0u_1u_2)\mu_1^-(u_2v_2v_0)\nu_0^-(v_3u_3u_2)\mu_2^+$, where the upper indices $\pm$ indicate either a forward or backward selection of orientation and each 2-path is presented with the orientation of the previously cited 5-cycle but must be presented in the next 5-cycle with its orientation reversed. Thus, Pet cannot be a $(\vec{C}_5; \vec{P}_3)$ -UH digraph.

For each positive integer n , let I_n stand for the n -cycle $(0, 1, \dots, n-1)$, where $0, 1, \dots, n-1$ are considered as vertices. If $G = Hea$ is the Heawood graph, then G can be obtained from I_{14} by adding the edges $(2x, 5+2x)$, for $x \in \{1, \dots, 7\}$ where operations are in $\mathbf{Z}_{14}$. The 28 6-cycles of G include the following 7 6-cycles: $\gamma_x = (2x, 1+2x, 2+2x, 3+2x, 4+2x, 5+2x)$, where $x \in \mathbf{Z}_7$. Then the following sequence of alternating 6-cycles and 3-arcs starts and ends up with opposite orientations for γ_0 :

$$\gamma_0^+(2345)\gamma_1^-(7654)\gamma_2^+(6789)\gamma_3^-(ba98)\gamma_4^+(abcd)\gamma_5^-(10dc)\gamma_6^+(0123)\gamma_0^-,$$

(where tridecimal notation is used, up to $d = 13$). Thus, Hea cannot be a $(\vec{C}_7; \vec{P}_4)$ -UH digraph.

If $G = Pap$ is the Pappus graph, then G can be obtained from I_{18} by adding the edges $(1+6x, 6+6x), (2+6x, 9+6x), (4+6x, 11+6x)$, for $x \in \{0, 1, 2\}$, where operations are mod 18. Then G admits a $(18\vec{C}_6; \vec{P}_3)$ -OAC formed by the oriented 6-cycles $A_0 = (123456), B_0 = (3210de), C_0 = (34bcde), D_0 = (0165gh), E_0 = (4329ab)$, (where octodecimal notation is used, up to $h = 17$), the 6-cycles A_x, B_x, C_x, D_x, E_x obtained by adding $6x$ mod 18 to (the integer representations of) the vertices of A_0, B_0, C_0, D_0, E_0 , where $x \in \mathbf{Z}_3 \setminus \{0\}$, and finally the 6-cycles $F_0 = (23ef89), F_1 = (hg54ba), F_2 = (61idc7)$.

If $G = Dod$ is the dodecahedral graph, then G can be seen as a 2-covering graph of the Petersen graph H , where each vertex u_x , (resp., v_x), of H is covered by two vertices a_x, c_x , (resp. b_x, d_x). This can be done so that a $(12\vec{C}_5; \vec{P}_2)$ -OAC of G is formed by the oriented 5-cycles $(a_0a_1a_2a_3a_4), (c_4c_3c_2c_1c_0)$ and for each $x \in \mathbf{Z}_5$ also $(a_xd_xb_{x-2}d_{x+1}a_{x+1})$ and $(d_xb_{x+2}c_{x+2}c_{x-2}b_{x-2})$.

If $G = Des$ is the Desargues graph, then G can be obtained from the 20-cycle I_{20} , with vertices $4x, 4x+1, 4x+2, 4x+3$ redenoted alternatively x_0, x_1, x_2, x_3 , respectively, for $x \in \{0, \dots, 4\}$, by adding the edges $(x_3, (x+2)_0)$ and $(x_1, (x+2)_2)$, where operations are mod 5. Then G admits a $(20\vec{C}_6; \vec{P}_3)$ -OAC formed by the oriented 6-cycles A_x, B_x, C_x, D_x , for $x \in \{0, \dots, 4\}$, where

$$\begin{aligned} A_x &= (x_0x_1x_2x_3(x+1)_0(x+4)_3), & B_x &= (x_1x_0(x+4)_3(x+4)_2(x+2)_1(x+2)_2), \\ C_x &= (x_2x_1x_0(x+3)_3(x+3)_2(x+3)_1), & D_x &= (x_0(x+4)_3(x+1)_0(x+1)_1(x+3)_2(x+3)_3). \end{aligned}$$

If $G = Cox$ is the Coxeter graph, then G can be obtained from the three 7-cycles $(u_1 u_2 u_3 u_4 u_5 u_6 u_0)$, $(v_1 v_3 v_5 v_0 v_2 v_4 v_6)$, $(t_1 t_4 t_0 t_3 t_6 t_2 t_5)$ by adding a copy of $K_{1,3}$ with degree-3 vertex z_x and degree-1 vertices u_x, v_x, t_x , for each $x \in \mathbf{Z}_7$. Then Cox admits the $(24\vec{C}_7; \vec{P}_3)$ -OAC

$$\begin{array}{lll} 0^1 = (u_1 u_2 u_3 u_4 u_5 u_6 u_0), & 0^2 = (v_1 v_3 v_5 v_0 v_2 v_4 v_6), & 0^3 = (t_1 t_4 t_0 t_3 t_6 t_2 t_5), \\ 1^1 = (u_1 z_1 v_1 v_3 z_3 u_3 u_2), & 1^2 = (z_4 v_4 v_2 v_0 z_0 t_0 t_4), & 1^3 = (t_6 t_2 t_5 z_5 u_5 u_6 z_6), \\ 2^1 = (v_5 z_5 u_5 u_4 u_3 z_3 v_3), & 2^2 = (t_6 z_6 v_6 v_4 v_2 z_2 t_2), & 2^3 = (u_1 z_1 t_1 t_4 z_0 u_0), \\ 3^1 = (v_5 v_0 z_0 u_0 u_6 u_5 z_5), & 3^2 = (z_4 t_4 t_1 z_1 v_1 v_6 v_4), & 3^3 = (t_6 t_2 z_2 u_2 u_3 z_3 t_3), \\ 4^1 = (u_1 u_0 z_0 v_0 v_2 z_2 u_2), & 4^2 = (t_6 t_3 z_3 v_3 v_1 v_6 z_6), & 4^3 = (z_4 u_4 u_5 z_5 t_5 t_1 t_4), \\ 5^1 = (z_4 u_4 u_3 u_2 z_2 v_2 v_4), & 5^2 = (v_5 v_3 v_1 z_1 t_1 t_5 z_5), & 5^3 = (t_6 z_6 u_6 u_0 z_0 t_0 t_3), \\ 6^1 = (z_4 v_4 v_6 z_6 u_6 u_5 u_4), & 6^2 = (v_5 v_3 z_3 t_3 t_0 z_0 v_0), & 6^3 = (u_1 u_2 z_2 t_2 t_5 t_1 z_1), \\ 7^1 = (u_1 u_0 u_6 z_6 v_6 v_1 z_1), & 7^2 = (v_5 z_5 t_5 t_2 z_2 v_2 v_0), & 7^3 = (z_4 t_4 t_0 t_3 z_3 u_3 u_4). \end{array}$$

If $G = Tut$ is Tutte's 8-cage, then G can be obtained from I_{30} , with vertices $5x, 5x+1, 5x+2, 5x+3, 5x+4, 5x+5$ denoted alternatively $x_0, x_1, x_2, x_3, x_4, x_5$, respectively, for $x \in \mathbf{Z}_5$, by adding the edges $(x_5, (x+2)_0)$, $(x_1, (x+1)_4)$ and $(x_2, (x+2)_3)$. Then G admits the $(90\vec{C}_8; \vec{P}_5)$ -OAC formed by the oriented 8-cycles

$$\begin{array}{lll} A^0 = (4_5 0_0 0_1 0_2 0_3 0_4 0_5 1_0), & B^0 = (4_2 4_3 4_4 4_5 1_0 1_1 1_2 1_3), & C^0 = (0_2 0_3 0_4 4_1 4_0 2_5 2_4 2_3), \\ D^0 = (3_3 3_2 3_1 4_4 4_3 4_2 1_3 1_2), & E^0 = (4_5 1_0 0_5 0_4 4_1 4_0 3_5 0_0), & F^0 = (4_5 0_0 3_5 4_0 2_5 2_4 1_1 1_0), \\ G^0 = (1_0 1_1 2_4 2_3 0_2 0_1 0_0 4_5), & H^0 = (2_3 2_4 1_1 1_0 0_5 0_4 0_3 0_2), & I^0 = (0_1 0_2 0_3 0_4 4_1 4_2 1_3 1_4), \\ J^0 = (1_0 0_5 0_4 0_3 3_2 3_1 4_4 4_5), & K^0 = (3_1 3_2 0_3 0_2 0_1 0_0 4_5 4_4), & L^0 = (2_3 2_4 2_5 3_0 3_1 3_2 0_3 0_2), \\ M^0 = (3_5 4_0 4_1 0_4 0_3 0_2 0_1 0_0), & N^0 = (3_5 3_4 2_1 2_0 1_5 1_4 0_1 0_0), & O^0 = (4_2 4_3 2_2 2_1 3_4 3_3 1_2 1_3), \\ P^0 = (4_5 4_4 4_3 4_2 4_1 0_4 0_5 1_0), & Q^0 = (4_0 4_1 4_2 1_3 1_4 1_5 3_0 2_5), & R^0 = (0_1 0_2 0_3 3_2 3_1 3_0 1_5 1_4), \end{array}$$

together with those obtained from these 18 8-cycles by adding $x \in \mathbf{Z}_5$ uniformly mod 5 to all subindices. Accordingly, these 8-cycles are denoted $A^x, \dots, R^x$, where $x \in \mathbf{Z}_5$.

If $G = Fos$ is the Foster graph, then G can be obtained from I_{90} , with vertices $5x, 5x+1, 5x+2, 5x+3, 5x+4, 5x+5$ denoted alternatively $x_0, x_1, x_2, x_3, x_4, x_5$, respectively, for $x \in \mathbf{Z}_{15}$, by adding the edges $(x_4, (x+2)_1)$, $(x_0, (x+2)_5)$ and $(x_2, (x+6)_3)$. The 90 10-cycles of G include the following 15 10-cycles, where $x \in \mathbf{Z}_{15}$.

$$\phi_x = (x_4 x_5 (x+1)_0 (x+1)_1 (x+1)_2 (x+1)_3 (x+1)_4 (x+1)_5 (x+2)_0 (x+2)_1),$$

Then the following sequence of alternating 10-cycles and 4-arcs:

$$\phi_0^+ [1_4] \phi_1^- [3_1] \phi_2^+ [3_4] \phi_3^- [5_1] \phi_4^+ [5_4] \phi_5^- [7_1] \phi_6^+ [7_4] \phi_7^- [9_1] \phi_8^+ [9_4] \phi_9^- [b_1] \phi_a^+ [b_4] \phi_b^- [d_1] \phi_c^+ [d_4] \phi_d^- [0_1] \phi_e^+ [0_4]$$

continues with ϕ_0^- , of opposite orientation to that of the initial ϕ_0^+ , where $[x_j]$ stands for a 3-path starting at the vertex x_j in the previously cited (to the left) oriented 10-cycle. Thus, Fos cannot be a $(\vec{C}_{10}; \vec{P}_5)$ -UH digraph.

The case of the Biggs-Smith graph is treated separately in [8]. $\square$

3 ‘Zipping’ the $(k-1)$ -powers of g -cycles ...

Given a CDT graph G with $\kappa = 3$, consider the collection $\mathcal{C}_g^{k-1}(G)$ of $(k-1)$ -powers of oriented g -cycles in the $(\eta\vec{C}_g; \vec{P}_k)$ -OAC of G in the proof of Theorem

2. If $k = 3$, then each arc $\vec{e}$ of a member C^2 of $\mathcal{C}_g^2(G)$ is marked by the middle vertex of the 2-arc $\vec{E}$ in C for which $\vec{e}$ stands, while the tail and head of $\vec{e}$ are marked by the tail and head of $\vec{E}$, respectively. This is the case in below and in [8]: we consider the CDT graphs G with $\kappa = k = 3$ in order to ‘zip’ such C^2 s along their O-O arc pairs to obtain corresponding graphs $Y(G)$ with $\mathcal{C}$ -UH properties. In all these cases, the following sequence of operations is performed:

$$G \rightarrow (\eta\vec{C}_g; \vec{P}_k)\text{-OAC}(G) \rightarrow \mathcal{C}_g^{k-1}(G) \rightarrow Y(G).$$

The CDT graphs G with $\kappa = 0$ do not admit the approach suggested in the previous paragraph for their g -cycles lack a $(\vec{C}_g; \vec{P}_k)$ -OA; those with $\kappa = 1$ admit the approach with $Y(G) = G$ so nothing new is obtained more than a corresponding polyhedral graph (embeddable into the sphere) with faces delimited by g -cycles, namely the tetrahedral, 3-cube and dodecahedral graphs; those with $\kappa = 2$ again admit the approach, but since $\kappa/2 = k - 1$, then $Y(G) = (g - 1)G^{k-1}$, the multigraph of multiplicity $g - 1$ on the $(k - 1)$ -th power of G .

If G is either the Pappus graph Pap or the Desargues graph Des , then $\mathcal{C}_6^2(G)$ is formed by triangles conforming a graph $Y(G)$ with just two connected components $Y_1(G)$ and $Y_2(G)$.




Figure 1: Toroidal cutouts of $Y_1(Pap)$ and $Y_2(Pap)$

Each of $Y_1(Pap)$ and $Y_2(Pap)$ is embeddable in a closed orientable surface T_1 of genus 1, or 1-torus. In fact, Figure 1 shows toroidal cutouts of $Y_1(Pap)$ and $Y_2(Pap)$. Notice that the copies of K_3 in $\mathcal{C}_6^2(G)$ are contractible in T_1 . These triangles form two collections $\mathcal{H}_1, \mathcal{H}_2$ of copies y_i^j of K_3 closed under parallel translation, where $y = A, B, C, D, E, F$; $i = 0, 1, 2$ and $j = 1, 2$, namely: the nine of $\mathcal{H}_1$ ($\mathcal{H}_2$) with horizontal edge below (above) its opposite vertex. There is also a collection $\mathcal{H}_0$ of nine non-contractible copies of K_3 in G , traceable linearly in three different parallel directions, three such triangles per direction, with the edges of each triangle marked by an associated common vertex of Pap .

There are embeddings of $Y_1(Pap)$ and $Y_2(Pap)$ in T_1 for which $\mathcal{H}_0$ and either $\mathcal{H}_1$ or $\mathcal{H}_2$ provide the composing faces. In addition, each of $\mathcal{H}_1$, $\mathcal{H}_2$ and $\mathcal{H}_0$ is formed by three classes of parallel elements, in the sense that any two of them in such a class do not have vertices in common. The self-dual (9_3) -configuration in the following theorem is the Pappus configuration, [5].

Theorem 3 $Y_1(Pap)$ and $Y_2(Pap)$ are isomorphic K_2 -fastened $\{H_0, H_1, H_2\}$ -UH graphs, where H_i is a representative of $\mathcal{H}_i$, for $i = 0, 1, 2$. Moreover, each of $Y_1(Pap)$ and $Y_2(Pap)$ can be taken as the Menger graph of the Pappus self-dual (9_3) -configuration in 12 different forms, by selecting the point set $\mathcal{P}$ and the line set $\mathcal{L} \neq \mathcal{P}$ so that $\{\mathcal{P}, \mathcal{L}\} \subset \{V(Pap), \mathcal{H}_0, \mathcal{H}_1, \mathcal{H}_2\}$ and the incidence relation either as the inclusion of a vertex in a copy of K_3 or as the containment by a copy of K_3 of a vertex or as the sharing of an edge by two copies of K_3 .

Proof. The statement can be established by managing the data given above. The 12 different claimed forms correspond to the arcs of the complete graph on vertex set $\{V(Pap), \mathcal{H}_0, \mathcal{H}_1, \mathcal{H}_2\}$. $\square$

If $G = Des$, then $Y_1(G)$ and $Y_2(G)$ are isomorphic K_2 -fastened (K_4, K_3) -UH graphs, each formed by five copies of K_4 and ten copies of K_3 , with each such copy of K_3 : **(a)** not forming part of a copy of K_4 in $Y_1(G)$ or $Y_2(G)$; **(b)** having its edges marked by a constant symbol, as shown in Figure 2.




Figure 2: Representations of $Y_1(Des)$ and $Y_2(Des)$

Deleting a copy H of K_4 from such $Y_i(Des)$ yields a copy of $K_{2,2,2}$, four of whose composing copies of K_3 , with no common edges, are faces of corresponding copies of $K_4 \neq H$; the other four copies of K_3 are among the ten mentioned copies of K_3 in G . A realization of $Y_1(G)$ (or $Y_2(G)$) in 3-space can be obtained from a regular octahedron O_3 realizing the $K_{2,2,2}$ cited above via the midpoints of the four segments joining the barycenters of four edge-disjoint alternate triangles in O_3 to the barycenter of O_3 by constructing the tetrahedra

determined by each of these alternate triangles and the nearest constructed mid-point, as well as the fifth central tetrahedron determined by the four midpoints.

By considering the barycenters of the resulting five tetrahedra and the segments joining them, a copy of K_5 in 3-space is obtained. The geometric line graph $L(K_5)$ it gives place to appears as a smaller version of $Y_1(G)$ (or $Y_2(G)$) contained in a octahedron $O'_3 \subset O_3$. This procedure may be repeated indefinitely, generating a sequence of realizations of $Y_1(G)$ (or $Y_2(G)$) in 3-space. Since $Y_1(Des)$ and $Y_2(Des)$ are isomorphic to $L(K_5)$, whose complement is Pet , then this sequence yields a corresponding sequence of realizations of Pet in 3-space.

We notice that the ten vertices and ten copies of K_3 of either $Y_i(Des)$ ($i = 1, 2$) may be considered as the points and lines of the Desargues self-dual (10_3) configuration, and that the Menger graph of this coincides with $Y_i(Des)$, [5]. Each vertex of $Y_i(Des)$ is the meeting vertex of two copies of K_4 and three copies of K_3 not forming part of a copy of K_4 .

Theorem 4 $Y_1(Des)$ and $Y_2(Des)$ are K_2 -fastened $\{K_4, K_3\}$ -UH graphs composed by five copies of K_4 and ten copies of K_3 each. Moreover, the ten vertices and ten copies of K_3 in either graph constitute the Desargues self-dual (10_3) configuration, which has the graph itself as its Menger graph. Furthermore, both graphs are isomorphic to $L(K_5)$, whose complement is the Petersen graph. $\square$

Theorem 4 can be partly generalized by replacing $L(K_5)$ by $L(K_n)$ ($n \geq 4$). This produces a K_2 -fastened $\{K_{n-1}, K_3\}$ -UH graph.

Theorem 5 The line graph $L(K_n)$, with $n \geq 4$, is a K_2 -fastened $\{K_{n-1}, K_3\}$ -UH graph with n copies of K_{n-1} and $\binom{n}{3}$ copies of K_3 .

Proof. We assume that each vertex of K_n is taken as a color of edges of $L(K_n)$ under the following rule: Color all the edges between vertices of $L(K_n)$ representing edges incident to a vertex v of K_n with color v . Then, each triple of edge colors for $L(K_n)$ corresponds to the edges of a well determined copy of K_3 in $L(K_n)$. Thus, there are exactly $\binom{n}{3}$ copies of K_3 intervening in $L(K_n)$ looked upon as a $\{K_{n-1}, K_3\}$ -UH graph. $\square$

The Fano plane $\mathcal{F}$, with point set $J_7 = \{1, \dots, 7\}$ and line set $\{124, 235, 346, 457, 561, 672, 713\}$, yields a coloring of the vertices and edges of $G = Cox$, represented on the upper left of Figure 3.

Observe that the colors of each vertex v of G and its three incident edges form a quadruple q whose complement $\mathcal{F} \setminus q$ is a triangle of $\mathcal{F}$ used as a ‘customary’ vertex denomination for v , [11], page 69. Then: **(a)** the triple formed by the colors of the edges incident to each vertex of G is a line of $\mathcal{F}$; **(b)** the color of each edge e of G together with the colors of the endvertices of e form a line of $\mathcal{F}$. In this representation of G , the vertices u_x, z_x, v_x, t_x in the proof of Theorem 2 are depicted concentrically from the outside to the inside, respectively, starting with $i = 1$, say, on the upper middle vertices with ‘customary’ vertex denominations 257, 134, 356, 567.

The squares C^2 of the 24 7-cycles C of G are taken with the following (cyclically presented) orientations, where each vertex v (resp. edge e) of a C^2 is indicated by the color of v (resp., subindicated by the color of the middle vertex of the 2-path of C that e represents). In fact, the resulting oriented 7-cycles can be denoted i^j , where $i \in \{0\} \cup J_7$ and $j \in J_3 = \{1, 2, 3\}$, namely:

0^1 :	(12345671234567) ;	0^2 :	(13572461357246) ;	0^3 :	(15263741526374) ;
1^1 :	(15467321546732) ;	1^2 :	(17534261753426) ;	1^3 :	(14725631472563) ;
2^1 :	(12543761254376) ;	2^2 :	(13275641327564) ;	2^3 :	(15362471536247) ;
3^1 :	(13476521347652) ;	3^2 :	(16354271635427) ;	3^3 :	(14623751462375) ;
4^1 :	(17435621743562) ;	4^2 :	(15764231576423) ;	4^3 :	(14527361452736) ;
5^1 :	(14326571432657) ;	5^2 :	(16453721645372) ;	5^3 :	(13674251367425) ;
6^1 :	(17236541723654) ;	6^2 :	(16752431675243) ;	6^3 :	(12647351264735) ;
7^1 :	(17632451763245) ;	7^2 :	(12746531274653) ;	7^3 :	(16257341625734) .




Figure 3: $\mathcal{F}$ -colored Cox and the three charts of the Klein graph $Y(Cox)$

Each 2-arc of G is cited exactly once in the oriented cycles i^j . Each 2-path of G appears twice in the i^j s, once for each one of its two 2-arcs. The assumed orientation of each C^2 corresponds with the orientation of the corresponding 7-cycle C . Thus, we consider each 7-cycle C with the orientation it induces in the corresponding C^2 , say i^j . In this case, we denote such a C as $\underline{i}^j$.

Each 2-path $\underline{e}$ of G separates two of the 24 7-cycles of G , say $\underline{i}^j$ and $\underline{k}^\ell$, with their orientations opposite over $\underline{e}$. Now, $\underline{i}^j$ and $\underline{k}^\ell$ restrict to the two

different 2-arcs provided by $\underline{e}$, say 2-arcs e_1 and e_2 . Then, e_1 and e_2 represent corresponding arcs $\underline{e}_1$ and $\underline{e}_2$ in i^j and k^ℓ , respectively.

Let us see that $\underline{e}_1$ and $\underline{e}_2$ can be ‘zipped’ into an edge e in the graph $Y(G)$. This, which happens to be the Klein graph, [1], graph 56B, can be assembled from the three charts shown on the upper right and bottom of Figure 3 by ‘zipping’ the 7-cycles i^j , interpreted all with counterclockwise orientation. Each of these three charts conforms a ‘rosette’, where the 7-cycles i^j with $i \neq 0$ are represented as ‘petals’ of the ‘central’ 7-cycles 0^1 , 0^2 and 0^3 . (The assemblage of $Y(G)$ can be done also around the 7-cycles i^1 , i^2 and i^3 , for any $i \in J_7$, not just $i = 0$). Moreover, each edge e in the external border of one of the three charts is accompanied by the denomination of a 7-cycle i_j incident to e which is a petal in one of the other two rosettes. Thus, the three charts can be assembled into the claimed graph $Y(G)$. Moreover, the 24 7-cycles i^j can be filled each with a corresponding 2-cell, so because of the cancelations of the two opposite arcs on each edge of $Y(G)$ (for having opposite orientations makes them mutually cancelable), $Y(G)$ becomes embedded into a closed orientable surface T_3 . As for the genus of T_3 , observe that

$$|V(Y(G))| = 2 \times 28 = 56 \text{ and } |E(Y(G))| = 2|E(G)| = 2 \times 42 = 84,$$

so that by the Euler characteristic formula for T_3 here,

$$|V(Y(G))| - |E(Y(G))| + |F_7(Y(G))| = 56 - 84 + 24 = -4 = 2 - 2.g(T),$$

and thus $g = 3$, so T_3 is a 3-torus. This yields the Klein map (of Coxeter notation) $\{7, 3\}_8$, (see [17]; note: the Petrie polygons of this map are 8-cycles).

Theorem 6 *The Klein graph $Y(Cox)$ is a $(C_7; P_2)$ -UH graph composed by 24 7-cycles that yield the Klein map $\{7, 3\}_8$ in T_3 . $\square$*

For the Klein map $\{7, 3\}_8$, the 3-torus appeared originally dressed as the Klein quartic $x^3y + y^3z + z^3x = 0$, a Riemann surface and the most symmetrical curve of genus 3 over the complex numbers. The automorphism group for this Klein map is $PSL(2, 7) = GL(3, 2)$, the same group as for $\mathcal{F}$, whose index is 2 in the automorphism groups of Hea , Cox and $Y(Cox)$.

Corollary 7 *The graph $Y'(Cox)$ whose vertices are the 7-cycles i_j of $Y(Cox)$, with adjacency between two vertices if their representative 7-cycles have an edge in common, is regular of degree 7, chromatic number 8 and has a natural triangular T_3 -embedding yielding the dual Klein map $\{3, 7\}_8$.*

Proof. Each vertex i_j of $Y'(Cox)$ is assigned color $i \in \{0\} \cup J_7$. Also, we have a partition of T_3 into 24 connected regions, each region having exactly seven neighboring regions, with eight colors needed for a proper map coloring. $\square$

References

- [1] I. Z. Bouwer et al., The Foster Census, R. M. Foster's Census of Connected Symmetric Trivalent Graphs, Charles Babbage Res. Ctr., Canada 1988.
- [2] A. E. Brouwer, A. M. Cohen and A. Neumaier, Distance-Regular Graphs, Springer-Verlag, New York 1989.
- [3] P. J. Cameron, *6-transitive graphs*, J. Combin. Theory Ser. B **28**(1980), 168–179.
- [4] G. L. Cherlin, The Classification of Countable Homogeneous Directed Graphs and Countable Homogeneous n -tournaments, Memoirs Amer. Math. Soc., vol. 131, number 612, Providence RI, January 1988.
- [5] H. S. M. Coxeter, *Self-dual configurations and regular graphs*, Bull. Amer. Math. Soc., **56**(1950), 413–455.
- [6] I. J. Dejter, *On a $\{K_4, K_{2,2,2}\}$ -ultrahomogeneous graph*, Australasian Journal of Combinatorics, **44**(2009), 63–76.
- [7] I. J. Dejter, *On a $\vec{C}_4$ -ultrahomogeneous oriented graph*, preprint 2009, accepted for publication in Discrete Mathematics.
- [8] I. J. Dejter, *On a $\{K_4, L(Q_3)\}_{K_3}$ -ultrahomogeneous graph*, manuscript 2010.
- [9] R. Fraïssé, *Sur l'extension aux relations de quelques propriétés des ordres*, Ann. Sci. École Norm. Sup. 71 (1954), 363–388.
- [10] A. Gardiner, *Homogeneous graphs*, J. Combinatorial Theory (B), **20** (1976), 94–102.
- [11] C. Godsil and G. Royle, Algebraic Graph Theory, Springer-Verlag, 2001.
- [12] Ja. Ju. Gol'fand and M. H. Klin, *On k -homogeneous graphs*, Algorithmic studies in combinatorics (Russian), bf 186(1978), 76–85.
- [13] D. C. Isaksen, C. Jankowski and S. Proctor, *On K_* -ultrahomogeneous graphs*, Ars Combinatoria, Volume LXXXII, (2007), 83–96.
- [14] A. H. Lachlan and R. Woodrow, *Countable ultrahomogeneous undirected graphs*, Trans. Amer. Math. Soc. 262 (1980), 51–94.
- [15] C. Ronse, *On homogeneous graphs*, J. London Math. Soc., **s2-17** (1978), 375–379.
- [16] J. Sheehan, *Smoothly embeddable subgraphs*, J. London Math. Soc., **s2-9** (1974), 212–218.
- [17] E. Schulte and J. M. Wills, *A Polyhedral Realization of Felix Klein's Map $\{3, 7\}_8$ on a Riemann Surface of Genus 3*, J. London Math. Soc., **s2-32** (1985), 539–547.